

# The prevalence of *Balantidium coli* in cattle and their breeders in some regions of Baghdad, Iraq

Asmaa Ghafer Hussin<sup>1</sup> Firas Rashad Al-Samarai<sup>2\*</sup>

<sup>1</sup>Al-Elam Sector for Primary Health Care/ Alkarkh Health Directorate Baghdad/ Ministry of Health/Iraq,

<sup>2</sup>Dept. Veterinary Public Health/Faculty of Veterinary Medicine/University of Baghdad/Iraq.

\*Corresponding author, E-mail address: [firmas\\_rashad@yahoo.com](mailto:firmas_rashad@yahoo.com)

## Abstract

A study was carried out to determine the prevalence of *Balantidium coli* infection in cattle and their breeders and related risk factors including the gender, age and region. A total of 200 cattle fecal samples was collected from different ages, sexes and regions of Baghdad (Hor-rageb, Abu Ghraib, AL-Jaderiya, AL-Makaseb, AL-Dowanem and AL-Hamdaniaa) and 88 breeder fecal samples from AL-Makaseb, AL-Dowanem and AL-Hamdaniaa. All these samples collected during the period from November - 2014 to May - 2015. The results showed that the prevalence of *Balantidium coli* were 29.50% and 9.09% in cattle and their breeders respectively. The association between infection and all studied factors was not significant except between infection and regions in cattle which were significant ( $P<0.01$ ). Based on the results of this research, it can be concluded that cattle are highly susceptible to balantidiasis irrespective of gender and age.

**Keywords:** *Balantidium coli*; cattle; cattle breeders; prevalence

## Introduction

In cattle, the disease problems, especially caused by protozoa lead to losses of productivity in terms of mortality, reduced milk, and meat (Roy et al., 2011). Therefore, it is necessary to identify the type of protozoa causing infection in the cattle to prevent, control and adopt the effective treatment. No doubt, the topic becomes more importance when the effect of protozoa is not limited to only cattle but extend to include humans also. Among the protozoan diseases, balantidiasis caused by *Balantidium coli* considered as one of an important disease that could infect animals and humans (Levine, 1995; Schuster and Ramirez-Avila, 2008). Several reports confirmed that, *B. coli* could be emerge as a significant pathogen that is able to cause disease in horses (Headley et al., 2008), buffaloes (Tarrar et al., 2008), cattle (Randhawa et al., 2010) and camels (Abubakr et al., 2000; Cox, 2005; AL-Tayib, 2014). However, the balantidiasis did not represent a serious problems in Japan, according to the conclusion demonstrated by Nakauchi, (1999) who found a low rate of infections in 56 mammalian species. Moreover, Thompson and Smith, (2011) stated that, although, *Balantidium* is known to cause infections in humans and is considered to be zoonotic but its role in zoonotic transmission is likely to be minimal.

The same situation exists in Islamic countries, although, balantidiasis is a zoonotic disease, a few researchers gave it their attention. This could be attributed to two reasons: Firstly, this disease can be found wherever pigs are found and in Islamic countries

like Iraq, pigs are not breeding because their religious beliefs prohibit consumption of pork. Secondly, *Balantidium* could infect pigs and other mammals along with humans, but its effect on the gastrointestinal tract usually not serious. It can thrive there in balance with its host without causing dysenteric symptoms, such as severe diarrhea and bloody stools. However, the problem could arise with the existence of malnutrition, alcoholism, or a compromised immune system virus [HIV]/AIDS, etc. that act to upset the balance in favor of the ciliate, leading to disease (Anargyrou et al., 2003; Vasilakopoulou et al., 2003; Ferry et al., 2004). On the other hand, there are other reasons can tip the balance in favor of estimate the prevalence of balantidiasis: according to Schuster and Ramirez-Avila, (2008), there are some fears of balantidiasis that can be expressed as the probability of increasing the prevalence of balantidiasis due to the warming of the earth's surface that could provide appropriate environment, particularly in the temperate areas of the world, for the survival of trophic and cystic stages of *Balantidium*. Moreover, Solaymani-Mohammadi and al., (2004) reported that wild boars were a reservoir for *Balantidium* in rural Western Iran, and it could be the source of infection in livestock and humans. The wild boars are existed in Iraq (Ministry of Environment, 2010), however, its role in infection was not studied. As, the water is the vehicle for most cases of balantidiasis, this disease could represent a problem in developing countries, as the opportunity of water sources to contaminate with porcine or human faeces is more likely to happen (Schuster and Ramirez-Avila, 2008). Bilal et al., (2009) stated that "There is considerable information on *B. coli* in most

pig raising communities in the world, but very little information on the parasite in cattle". Moreover, the information available is scanty about the prevalence of *Balantidium coli* in cattle and their breeders. Hence, this study was aimed to investigate the prevalence of *Balantidium coli* in cattle and their breeders along with identifying the association infection between them.

### Materials and Method

A total of 200 cattle fecal samples were collected from different ages, sexes and regions of Baghdad (Hor-rageb, Abu Ghraib, AL-Jaderiya, AL-Makaseb, AL-Dowanem and AL-Hamdaniaa) and 88 breeder fecal samples collected from AL-Makaseb, AL-Dowanem and AL-Hamdan.

All samples collected during the period from November- 2014 to May -2015. The faecal samples were collected directly from the rectum, in a clean plastic containers (100ml size) and were tightly closed and given sequential numbers. All information for the animal and breeder included age, sex and the name of region were recorded on a special form bearing the number of the sample, then the samples were transported in refrigerated bag to the zoonotic unite at the College of Veterinary Medicine in Baghdad University. After that, a portion of stool was examined at field by direct wet mount, Lugol's iodine staining technique and modified Ziehle-Nelsen staining to identify the *Balantidium coli*.

### Results and Discussion

Results revealed that the prevalence of *Balantidium coli* in cattle from different regions of Baghdad, Iraq was 29.50% (Table 1). The present estimation is higher than 27.5% in India (Niphadkar and Raote, 1994), 25.29% in Bangladesh (Rahman and Samad, 2010), 25% in Pakistan (Bilal et al., 2009), 17.19% in Indonesia (Wisesa et al., 2015) and 6.6% in Kenya (Kanyari et al., 2010) and lower than 45.45% in India (Palanivel et al. 2005) and 45.03% in Bangladesh (Roy et al., 2011). The variation in estimations of prevalence could be attributed to several factors such as breed, sample size, selection of samples, the technique of sample examination, agro-ecological, Geo-climatic conditions, management and nutrition (Hussin, 2015).

Table 1 showed the infection rates in male and female cattle were 32.30% and 28.14% respectively. The association between gender and infection was not significant. The result is consistent with the result obtained by Bilal et al., (2009), Kanyari et al., (2010) and Singh et al., (2012) in cattle. However, the result of this study contrasted with the result obtained by Wisesa et al., (2015) who found that the infection rate of balantidiasis in cattle was significantly ( $p<0.05$ ) higher in male (23.20%) than female (13.72%). On the other hand, opposite results were found by Bachal et al., (2002) who reported the infection rate of *Balantidium coli* in the female buffalo (48.30%) was higher than the male buffalo (45.12%). Roy et al., (2011) found the similar result and reported a higher prevalence of balantidiasis in female buffalo (47.32%) compared to male buffalo (38.46%).

The inconsistencies in the results could be belonged to the differences in the physiological condition of cattle ( pregnancy, lactation and parturition and lower feed supplements for production) which may cast a shadow on infection rate and lead to decreasing the resistance of the animal (Roy et al., 2011). The levels of prolactin and progesterone hormones affect the female. High levels of these hormones make female more susceptible to any infection (Lloyd, 1983). Meanwhile, male cattle are also susceptible to stress and this will decrease its resistance to infection (Mamun et al., 2011).

Concerning the difference among age groups, results showed that the differences were not significant (Table 2). These results supported the previous results obtained by Kanyari et al., (2010) and Wisesa et al., (2015). Meanwhile, our results disagreed with the results of Roy et al., (2011) who reported that age was significantly influenced the infection rate of balantidiasis.

The effect of regions on the infection rate of balantidiasis of cattle in the present study was significant ( $P < 0.01$ ) (Table 3). No infection of *Balantidium* was found in two regions (Hor-ragab and Al-Jaderia) while the highest infection rate was found in Al-Dowanem (56.75%). Similar results reported by Kanyari et al., (2010) and Wisesa et al., (2015). These differences could be attributed to variation in herd size, physiological status of animal, types of food and water, food supply systems, hygienic conditions, the level of veterinary care, and drugs or vaccinations (Hussin, 2015).

The results shown in (Table 4) showed that the prevalence of *Balantidium coli* in breeders (9.09%) was higher than 1.4% in Thailand (Chavalittamrong and Jirapinyo, 1984), 2.4% in children of India (Kaur et al., 2002) and 6.67% in Bangladesh (Al-Hasan et al., 2015). Although there are several factors could be responsible for these differences, but contact with animals could play a big role for these variations. The infection rates in male and female breeders were 12.12% and 7.20% respectively. However, the association between gender and infection was not significant. The present finding disagreed with the results obtained by Bui et al. (2008) who reported that the prevalence was higher in male (11.5%) than female (3.4%).

The non-significant effect of gender on the rate of infection is anticipated because, in rural regions, both males and females contact with cattle and exposed to the same pathogenic agents (Hussin, 2015). Despite the infection rate showed a decreasing trend with advanced age, but the association between age category and infection was not significant. (Table 5). These results disagreed with the results reported by Bui et al. (2008) and Al-Hasan et al., (2015) who confirmed the significant effect of age on infection rate of *Balantidium coli*.

The results also revealed that the association between infection and regions was not significant (Table 6). Based on the results of this research, it can be concluded that the high prevalence of *Balantidium coli* infection in cattle (29.50%) indicated that cattle are highly susceptible to balantidiasis irrespective of gender and age. On the other hand, the prevalence of

**Table 1:** Infection rate of *B. coli* according to gender in cattle

Gender	Total No.	+ve No	Percentage %
Male	65	21	32.30
Female	135	38	28.14
Chi-square value			0.37
P			0.55
Total	200	59	29.50

+ve = Positive cases

P = Probability

**Table 2:** Infection rate of *B. coli* according to age in cattle

Age/year	No.	+ve No.	Percentage %
<1	58	17	29.31
1-<3	49	13	26.53
3-<6	47	13	27.65
6≤	46	16	34.78
Chi-square value			0.90
P			0.82

+ve = Positive cases

P = Probability

**Table 3:** Infection rate of *B. coli* according to regions in cattle

Regions	No.	+ve No.	Percentage %
Hor-Rageb	31	0	0
Abu- Ghraib	34	7	20.58
AL-Jaderiya	26	0	0
AL-Makaseb	38	13	34.21
AL-Dowanem	37	21	56.75
AL-Hamданияa	34	18	52.94
Chi sq			47.75
P			<0.01

+ve = Positive cases

P = Probability

**Table 4:** Infection rate of *B. coli* according to gender in breeders

Gender	Total No.	+ve No	Percentage %
Male	33	4	12.12
Female	55	4	7.2
Chi-square value			0.58
P			0.44
Total	88	8	9.09

+ve = Positive cases

P = Probability

**Table 5:** Infection rate of *B. coli* according to age in breeders

Age/year	No.	+ve No.	Percentage %
<10	38	5	13.15
10-<30	27	2	7.40
30≤	23	1	4.34
Chi-square value			1.47
P			0.47

+ve = Positive cases

P = Probability

**Table 6:** Infection rate of *B. coli* according to age in breeders

Age/year	No.	+ve No.	Percentage %
AL-Makaseb	27	3	11.11
AL-Dowanem	32	4	12.50
AL-Hamdaniaa	29	1	3.44
Chi-square value			1.70
P			0.42

+ve = Positive cases

P = Probability

*Balantidium coli* in breeders is also high (9.09%), that will double the adverse effect of infection because it is known, that the presence of pathogenic bacteria (e.g., Salmonella) in the intestine can worsen an infection by invading colonic lesions caused by *Balantidium coli* (Levine, 1961; Skotarczak, 1997a, 1997b; Sestak et al., 2003). The results also reflect the poor sanitary condition, drinking of contaminated water and lack of knowledge about health and hygiene.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

The work was financially supported by the Faculty of Veterinary Medicine/University of Baghdad, Iraq.

### References

- Abubakr M, Nayel MN, Fadlalla M, Abdelrahman AO, Abuobeida S, Elgabara YM. Prevalence of gastrointestinal parasites in young camels in Bahrain. *Rev Elev Med Vet Pays Trop.* 2000; 53: 267-271.
- Al-Hasan A, Ali A, Al-Hasan A, Rakib FK, Alam MA, MM Mondal. Prevalence of *Balantidium coli* infection in man in Mymensingh Bangladesh *Int J Nat Soc Sci.* 2015; 2:33-36
- AL-Tayib O. Zoonotic balantidiasis in camel from Saudi Arabia. *Sch Acad J Biosci.* 2014; 2(7): 445-447
- Anargyrou KGL, Petrikkos MTE, Suller A, Skiada MR, Siakantaris RT, Osuntoyinbo G, Pangalis C, and Vaiopoulos G. Pulmonary *Balantidium coli* infection in a leukemic patient *Am J Hematol.* 2003; 73:180-183.
- Bachal B, Sharif P, Rahamatullah R, and Aijaz HS. Prevalence of gastro-intestinal helminths in Buffalo calves *Journal of Biological Sciences.* 2002; 2(1) 43-45.
- Bilal CQ, Khan MS, Avais M, Ijaz M, Khan JA. Prevalence and chemotherapy of *Balantidium coli* in cattle in the River Ravi region Lahore (Pakistan) *Veterinary Parasitology.* 2009; 163 : 1-2 7:15-17.
- Biu AA, and Dauda M. Prevalence study on enteric protozoans responsible for a copromicroscopical investigation in Sardinia *Parassitologia.* 2008; 48 (14): 313-314.
- Chavalittamrong B, Jirapinyo P. Intestinal parasites in pediatric patients with diarrhael diseases in Bangkok *Southeast Asian J Trop Med Public Health.* 1984; 15: 385-8.

- Cox FE. Human balantidiasis in Iran: are camels' reservoir hosts *Tred Parasitol.* 2005; 21: 553
- Ferry TD, Bouhour F, De Monbrison F, Laurent H, Domouchel-Champagne S, Picot M A, and P Granier. Severe peritonitis due to *Balantidium coli* acquired in France *Eur J Clin Microbiol Infect Dis.* 2004; 23:393-395.
- Headley SA, Kummala E, Sukura A. *Balantidium coli* infection in a Finnish horse *Vet Parasitol.* 2008; 158: 129-132.
- Hussin AG. Evaluation of different diagnostic methods for zoonotic intestinal protozoz in cattle camels and their breeders A thesis Faculty of Veterinary Medicine University of Baghdad Iraq. 2015.
- Kanyari PWN, Kagira JM, and Mhoma JRL. Prevalence of endoparasites in cattle within urban and peri-urban areas of Lake Victoria Basin Kenya with special reference to zoonotic potential *Sci Parasitol.* 2010; 11(4):171-178.
- Kaur R, Rawat D, Kakkar M, Uppal B, and Sharma VK. Intestinal parasites in children with diarrhea in Delhi India *Southeast Asian J Trop Med public Health.* 2002; 33(4):725-729.
- Levine ND. Protozoan parasites of Domestic Animals and of Man. Burgess Publishing Co., Minneapolis. 1961; Pp. 133.
- Levine ND. *Veterinary Protozoology* Ames: Iowa State University Press. 1995.
- Lloyd S. Effect of pregnancy and lactation up on infection. *Veterianry Immunology Immunopathology.* 1983; 4: 153-176.
- Mamun MAA, Begum N, and Mondal MMH. A coprological survey of gastro-intestinal parasites of water buffaloes (*Bubalus bubalis*) in Kurigram district of Bangladesh. *J Bangladesh Agril Univ.* 2011; 9(1): 103-109.
- Ministry of Environment. Iraqi fourth national report to the convention on biological diversity Iraq. 2010.
- Nakauchi K. The prevalence of *Balantidium coli* infection in fifty-six mammalian species *J Vet Med Sci.* 1999; 61:63-65.
- Palanivel KM, Thangathurai R, and Nedunchellian S. Epizootiology of *Balantidium coli* infection in ruminants. *Indian Veterinary Journal.* 2005; 82(6): 682-683.
- Rahman MM and Samad MA. Prevalence of sub-clinical gastro-intestinal parasitosis and their effects on milk production with therapeutic management in Red Chittagong cattle. *Bangladesh Journal of Veterinary Medicine.* 2010; 8: 11-16.

- Randhawa S, Singla LD, Randhawa C. Chronic cattle diarrhea due to *Balantidium coli* infection-a clinical report J Vet Parasitol. 2010; 24:197-198.
- Roy BC, Mondal MMH, Talukder MH, and Majumder S. Prevalence of *Balantidium coli* in buffaloes at different areas of Mymensingh. J Bangladesh Agr Univ. 2011; 9: 67-72.
- Schuster FL and Ramirez-Avila L. Current world status of *Balantidium coli*. Clinical Microbiology Reviews. 2008; 21(4):626-638.
- Sestak KCK, Merritt J, Borda E, Saylor SR, Schwamberger F, Cogswell ES, Didier PJ, Didier G, Plauche RP, Bohn PP, Aye P, Alexa RL and Lackner AA. Infectious agent and immune response characteristics of chronic enterocolitis in captive rhesus macaques. Infect Immun. 2003; 71: 4079-4086.
- Singh NK, Singh H, Haque M and Rath SS. Prevalence of parasitic infections in buffaloes in and around Ludhiana District Punjab India: A Preliminary Study. J Parasit Dis. 2012; 36(2): 256-259.
- Skotarczak B. Bacterial-flora in acute and symptom-free balantidiasis Acta Parasitol. 1997a; 42: 230-233.
- Skotarczak B. 1997b. Ultrastructural and cytochemical identification of peroxisomes in *Balantidium coli* Ciliophora Folia Biol 45:117-120
- Solaymani-Mohammadi S, Rezaian M, Hooshyar H, Mowlavi GR, Babaei Z, Anwar MA. Intestinal protozoa in wild boars (*Sus scrofa*) in Western Iran. J Wildl Dis. 2004; 40: 801-803.
- Tarrar MA, Khan MS, Pervez K, Ashraf K, Khan JA and Rehman ZU. Detection and chemotherapy of *Balantidium coli* in buffaloes around Lahoure. Pakistan. Pak J Agri Sci. 2008; 45(2): 163-166.
- Thompson RCA and Smith A. Zoonotic enteric protozoa. Veterinary Parasitology. 2011; 182: 70-78.
- Vasilakopolou AK, Dimarongona A, Samakovli K, Papadimitris and Avlami A. *Balantidium coli* pneumonia in an immunocompromised patient. Scand J Infect Dis. 2003; 35: 144-146.
- Wisasa IBGR, Siswanto FM, Putra TA, Oka IBM, Suratma NA. Prevalence of *Balantidium* sp in Bali cattle at different areas of Bali. International Journal of Agriculture Forestry and Plantation. 2015; 1:49-53.