



Original Article

Phylogenetic relationships of *Kaempferia* plants based on inter-simple sequence repeat fingerprints

Orawan Theanphong^{1*}, Thaya Jenjittikul², Withawat Mingvanish³,
and Kanchana Rungsihirunrat⁴

¹ Department of Pharmacognosy, Faculty of Pharmacy, Rangsit University,
Mueang, Pathum Thani, 12000 Thailand

² Department of Plant Science, Faculty of Science, Mahidol University,
Ratchathewi, Bangkok, 10400 Thailand

³ Department of Chemistry, Faculty of Science, King Mongkut's University of Technology Thonburi,
Thung Khru, Bangkok, 10140 Thailand

⁴ College of Public Health Sciences, Chulalongkorn University,
Pathum Wan, Bangkok, 10330 Thailand

Received: 29 July 2016; Revised: 30 October 2016; Accepted: 8 March 2017

Abstract

Plants in genus *Kaempferia* (Zingiberaceae) have been used as folk medicine in several countries. Identification and classification of *Kaempferia* plants are difficult because of the morphological similarity between species. In this study, phylogenetic relationships of *Kaempferia* species were employed by inter-simple sequence repeat (ISSR) fingerprints. Ten ISSR primers of 45 ISSR primers produced 186 bands with an average 19 bands per primer. The similarity index (SI) ranged from 0.0023-0.9950. A dendrogram was constructed using unweighted pair-group method of the arithmetic average (UPGMA). The results showed that the *Kaempferia* species could be divided into two main groups. The phylogenetic relationships were associated with the morphological characteristics.

Keywords: *Kaempferia*, Zingiberaceae, inter-simple sequence repeat fingerprints

1. Introduction

The genus *Kaempferia* is a medium-sized rhizomatous herb belonging to the Zingiberaceae family. *Kaempferia* is comprised of approximately 40 species distributed in monsoonal tropical Asia (Skornickova & Newman, 2015), of which 29 are found in Thailand (Pooma & Suddee, 2014). In Thailand, *Kaempferia* plants are ethnomedically used in

the treatment of flatulence, fever, gastric ulcer, leucorrhoea, oedema, and healing of wounds (Chuakul, 2005; Chuakul & Boonpleng, 2003, 2004; Maneenoon *et al.*, 2015; Nuammee *et al.*, 2012; Suksri *et al.*, 2005; Tangjitman *et al.*, 2015).

However, the taxonomic identification and classification of these plants based on morphological characteristics is not always satisfactory because of their similar appearance within the genus and also with other closely related genera, i.e. *Boesenbergia*, *Scaphochlamys*, *Caulokaempferia*, and *Curcuma* (Kitamura *et al.*, 2007; Picheansoonthon & Koonterm, 2008; Syamkumar & Sasikumar, 2007; Techaprasan *et al.*, 2010; Vanijajiva *et al.*, 2005; Zou *et al.*, 2011).

*Corresponding author
Email address: theanphong@gmail.com

Nowadays, molecular markers are valuable tools in the characterization and evaluation of genetic diversity within and between species and populations (Heubl, 2010; Semagn *et al.*, 2006; Tharachand *et al.*, 2012). The inter-simple sequence repeat (ISSR) technique is a fast and inexpensive technique for the characterization of genetic relatedness among populations (Dogan *et al.*, 2016; Prevost & Wilkinson, 1999). It is the marker of choice for the amplification of variations of genomic sequences between microsatellites using a single primer of nucleotide repeat units. ISSR employs simple sequence repeats in the primers so that it does not require sequence information for primer design, which is a major advantage of ISSR analysis (Son *et al.*, 2012). The ISSR marker has been used in phylogenetic analysis in several plants such as *Allium* (Hao *et al.*, 2002; Son *et al.*, 2012), *Oryza* (Joshi *et al.*, 2000); *Phyllanthus* (Rout & Aparajita, 2010), *Matthiola* (Dagan *et al.*, 2016), and *Klasea* (Dogan *et al.*, 2015). However, the number of studies of the genetic relationships of *Kaempferia* species using the ISSR marker is quite limited.

Thus, the aims of this study were to investigate the genetic diversity and phylogenetic relationships of *Kaempferia* species on the basis of ISSR fingerprinting analysis.

2. Materials and Methods

2.1 Plant materials

Fresh rhizomes of 11 accessions of *Kaempferia* belonging to 10 species were collected between May and July 2015 from different locations (9 accessions from Thailand and 2 accessions from Laos). Four Zingiberaceae plants; *Curcuma aromatica*, *Boesenbergia rotunda*, *Zingiber montanum* (tribe Zingibereae), *Alpinia galanga* (tribe Alpinieae), and *Citrus hystrix* (Rutaceae) were used as outgroup plants in this study. The details of plant samples used in the study are shown in Table 1. The plant samples

Table 1. Details of plant samples used in this study.

| Plant samples | Locality | Code |
|---|--------------|------|
| <i>Kaempferia angustifolia</i> Roscoe | Prachin Buri | KA |
| <i>K. elegans</i> (Wall.) Baker | Prachin Buri | KE |
| <i>K. galanga</i> L. | Chiang Mai | KG |
| <i>K. laotica</i> Gagnep. | Laos | KL |
| <i>K. larsenii</i> Siriruga | Ubon | KN |
| | Ratchathani | |
| <i>K. marginata</i> Carey ex Roscoe | Prachin Buri | KM |
| <i>K. parviflora</i> Wall. ex Baker | Prachin Buri | KP |
| (cultivar type) | | |
| <i>K. parviflora</i> Wall. ex Baker | Laos | KP2 |
| (wild type) | | |
| <i>K. pulchra</i> Ridl. | Kanchanaburi | KC |
| <i>K. rotunda</i> L. | Ratchaburi | KR |
| <i>K. siamensis</i> Siriruga | Sakon Nakhon | KS |
| <i>Curcuma aromatica</i> Salisb. | Chiang Mai | CA |
| <i>Zingiber montanum</i> (J.Koenig) | Bangkok | ZM |
| Link ex A.Dietr. | | |
| <i>Alpinia galanga</i> (L.) Willd. | Bangkok | AG |
| <i>Boesenbergia rotunda</i> (L.) Mansf. | Bangkok | BR |
| <i>Citrus hystrix</i> DC. | Pathum Thani | CH |

were authenticated by Assist. Prof. Dr. Thaya Jenjittikul (Department of Plant Science, Faculty of Science, Mahidol University, Thailand). The voucher specimens of these plants have been deposited at Department of Pharmacognosy, Faculty of Pharmacy, Rangsit University, Thailand. All plant samples were employed for cultivation at the Department of Pharmacognosy, Faculty of Pharmacy, Rangsit University, Bangkok, Thailand, for 4-6 weeks. Morphological characters of the cultivated plants were recorded, i.e. leaf position, leaf shape, leaf base, leaf apex, and position of inflorescence.

2.2 DNA isolation

Fresh young leaf of each plant was ground in liquid nitrogen with mortar and pestle to obtain a fine powder. Total genomic DNA was isolated from the fine powder using the DNeasy Plant Mini Kit (Qiagen, Germany) according to the manufacturer's protocol. DNA quantifications were checked by 1.5% agarose gel electrophoresis. Genomic DNA was kept at -20 °C for ISSR analysis.

2.3 Inter-simple sequence repeat (ISSR) amplification

The ISSR reaction was carried out in 25 µl containing 2 µl of genomic DNA, 1X amplification buffer, 5 mM MgCl₂, 0.4 mM of each dNTP, 1 U of *Taq* DNA polymerase (Fermentas, Canada), and 0.4 µM ISSR primers (Eurofins MWG Operon, Germany). The amplification was performed using a DNA thermal cycler (Applied Biosystems, USA) with an initial pre-denaturation at 95 °C for 1 min, denaturation at 95 °C for 45 sec, annealing at 45 °C for 45 sec, extension at 72 °C for 1 min with 35 cycles and final extension at 72 °C for 5 min. The ISSR products were separated alongside a molecular weight marker (1 kb Gene-Ruler, Fermentas, Canada) on 1.5% agarose gel in TBE buffer and stained with ethidium bromide. The ISSR fragments were photographed using a UV transilluminator and analyzed with a gel documentation system (Syngene, USA). The PCR amplifications were repeated at least three times in the same condition.

2.4 ISSR data analysis

The ISSR bands were scored as 0 or 1 for the absence or presence of bands, respectively. Only clear and reproducible bands were scored as 1. The similarity index was calculated from the data generated using Dice similarity index coefficient (Nei & Li, 1979). The dendrogram was constructed based on the similarity matrix data using the unweighted pair group method with arithmetic averages (UPGMA), clustering by FreeTree and TreeView software with 1000 bootstrap (Pavlicek *et al.*, 1999).

3. Results and Discussion

A total of 45 ISSR primers were used for initial screening; 10 primers produced 186 clear and reproducible bands in all plant samples. The total number of amplified products ranged from 12-24 bands with an average of 19 bands by each primer. The highest number of ISSR bands (24 bands) was generated from ISSR-38 primer while the lowest

number of bands (12 bands) was from ISSR-40. The approximate range of band size was 223-2505 bp. The percentage of polymorphism ranged from 9.25% to 41.67% as demonstrated by ISSR-27 and ISSR-40, respectively (Table 2). Change in the amplified products using ISSR primer can arise through the insertions/deletions or the loss of primer binding sites which generated different banding patterns.

Table 2. The sequence of ISSR primers and the number of PCR products obtained from plant samples.

| Primer | Nucleotide sequence (5' to 3') | No. of bands | Size of bands (bp) | Polymorphism (%) |
|---------|--------------------------------|--------------|--------------------|------------------|
| ISSR 02 | AGA GAG AGA GAG AGC | 18 | 252-1456 | 16.67 |
| ISSR 08 | GTG TGT GTG TGT GTC | 18 | 304-2505 | 33.33 |
| ISSR 19 | GAG AGA GAG AGA GAG GC | 22 | 264-2472 | 13.64 |
| ISSR 27 | AGA GAG AGA GAG AGA CYT | 21 | 259-1903 | 9.25 |
| ISSR 28 | AGA GAG AGA GAG AGA GYA | 17 | 237-1586 | 17.65 |
| ISSR 37 | TGT GTG TGT GTG TGT GRC | 16 | 267-1661 | 25.00 |
| ISSR 38 | AGC AGC AGC AGC AGC AGC | 24 | 223-1322 | 25.00 |
| ISSR 39 | ATG ATG ATG ATG ATG ATG | 22 | 242-2108 | 22.73 |
| ISSR 40 | CTC CTC CTC CTC CTC CTC | 12 | 278-1676 | 41.67 |
| ISSR 43 | GGG TGG GGT GGG GGT | 17 | 284-1727 | 23.53 |
| | Total | 186 | 223-2505 | |

According to 10 primers that produced clear and reproducible bands, the ISSR-27 primer generated the approximately 1000 bp characteristic band of all *Kaempferia*

species and four Zingiberaceous outgroup plants, this band was not observed in *C. hystrix* (Figure 1). Therefore, this band may be used as the characteristic band for Zingiberaceous plants.




Figure 1. ISSR fingerprint of 11 *Kaempferia* and outgroup plants obtained from the ISSR-27 primer. Abbreviations of the plant samples are according to codes used in Table 1. M: GeneRuler 1 kb (size shown in bp).

Since ISSR are dominant markers, the amplified bands are scored as diallelic. The pair-wise comparisons of the ISSR profiles based on both the shared and unique amplification bands were used to generate a similarity index. Among 11 accessions of *Kaempferia* plants including 5 outgroup plants, Dice similarity index ranged from 0.0023 to 0.9950 (Table 3). The highest genetic similarity index (0.9950) was found between *K. parviflora* cultivar and wild type, whereas the lowest (0.0023) was found between *K. elegans* and *Citrus hystrix*.

Table 3. Similarity matrix of *Kaempferia* and outgroup plants generated using Dice similarity coefficient.

| | KG | KM | KS | KC | KL | KP1 | KP2 | KE | KR | KA | KN | ZM | AG | CA | BR | CH |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| KG | 1.0000 | | | | | | | | | | | | | | | |
| KM | 0.9793 | 1.0000 | | | | | | | | | | | | | | |
| KS | 0.9170 | 0.9397 | 1.0000 | | | | | | | | | | | | | |
| KC | 0.8251 | 0.8053 | 0.8120 | 1.0000 | | | | | | | | | | | | |
| KL | 0.8957 | 0.8755 | 0.8880 | 0.9106 | 1.0000 | | | | | | | | | | | |
| KP1 | 0.4585 | 0.4688 | 0.4924 | 0.5892 | 0.5359 | 1.0000 | | | | | | | | | | |
| KP2 | 0.4622 | 0.4646 | 0.4886 | 0.5938 | 0.5399 | 0.9950 | 1.0000 | | | | | | | | | |
| KE | 0.4603 | 0.4706 | 0.4867 | 0.5914 | 0.5379 | 0.9756 | 0.9684 | 1.0000 | | | | | | | | |
| KR | 0.6015 | 0.6022 | 0.6202 | 0.6619 | 0.6528 | 0.7653 | 0.7573 | 0.7613 | 1.0000 | | | | | | | |
| KA | 0.6087 | 0.6093 | 0.6272 | 0.6690 | 0.6597 | 0.7588 | 0.7508 | 0.7548 | 0.9920 | 1.0000 | | | | | | |
| KN | 0.6081 | 0.6087 | 0.6127 | 0.6547 | 0.6456 | 0.7468 | 0.7386 | 0.7492 | 0.9849 | 0.9849 | 1.0000 | | | | | |
| ZM | 0.1504 | 0.1747 | 0.1772 | 0.1732 | 0.1765 | 0.2222 | 0.2162 | 0.2154 | 0.2324 | 0.2324 | 0.2278 | 1.0000 | | | | |
| AG | 0.0566 | 0.0644 | 0.0717 | 0.0669 | 0.0716 | 0.0978 | 0.0955 | 0.0928 | 0.1050 | 0.1075 | 0.1029 | 0.1947 | 1.0000 | | | |
| CA | 0.2097 | 0.2197 | 0.2275 | 0.2115 | 0.2308 | 0.2531 | 0.2504 | 0.2500 | 0.2838 | 0.2838 | 0.2750 | 0.2066 | 0.6011 | 1.0000 | | |
| BR | 0.2161 | 0.2222 | 0.2262 | 0.2251 | 0.2258 | 0.2547 | 0.2487 | 0.2517 | 0.2782 | 0.2782 | 0.2762 | 0.2202 | 0.6075 | 0.5035 | 1.0000 | |
| CH | 0.0024 | 0.0100 | 0.0032 | 0.0167 | 0.0098 | 0.0170 | 0.0076 | 0.0023 | 0.0514 | 0.0079 | 0.0947 | 0.0812 | 0.0694 | 0.0326 | 0.0212 | 1.0000 |

A dendrogram was constructed according to the UPGMA cluster analysis using the Dice similarity coefficient. Based on the dendrogram, *Kaempferia* species were categorized into two major groups. Cluster I and II were divided into two subgroups Ia/ Ib and IIa/IIb, respectively. Subgroup Ia, consisted of *K. elegans* and *K. parviflora* (cultivar and wild type), with a similarity index range of

0.9684-0.9950. Subgroup Ib included *K. larsenii*, *K. rotunda*, and *K. angustifolia*, with a similarity index range of 0.9849-0.9920. *K. siamensis*, *K. galangal*, and *K. marginata* were clustered into subgroup IIa with a similarity index range of 0.9170-0.9793, whereas *K. pulchra* and *K. laotica* were clustered into subgroup IIb with a 0.9106 similarity index. The separation of all subgroups and two major groups

received 100% bootstrap support. Four Zingiberaceous outgroup plants, *Zingiber montanum*, *Curcuma aromatica*, *Boesenbergia rotunda*, and *Alpinia galanga*, were completely separated from the *Kaempferia* species with 100% bootstrap support. In addition, *Citrus hystrix*, a dicotyledon plant, was completely separated from all Zingiberaceous plants with 100% bootstrap support (Figure 2). The results were similar to those previously reported. Based on ISSR fingerprints, *K.*

angustifolia and *K. rotunda* were clustered into the same group (Siriluck *et al.*, 2014) while based on RAPD fingerprints and chloroplast DNA sequences, *K. marginata* and *K. galanga* were clustered together (Techaprasan *et al.*, 2010; Theanphong *et al.*, 2013). In addition, *K. parviflora* and *K. elegans* were clustered into the same group on the basis of chloroplast DNA sequences (Techaprasan *et al.*, 2010).




Figure 2. Dendrogram produced by UPGMA cluster analysis of ISSR data showing the genetic relationship among 11 *Kaempferia* plants and outgroup plant.

The results obtained from this study were also correlated with the morphological characteristics. *Kaempferia* plants in cluster I were 6-30 cm tall and they had erect elliptic oblong leaves and inflorescence rising between the two innermost leaf-sheaths or rising from the rhizome together with a 1-10 cm petiole length. *K. siamensis*, *K. galanga*, *K. marginata*, *K. pulchra*, and *K. laotica* which were clustered into the second group were 2-10 cm tall and had horizontal elliptic-suborbicular leaves and inflorescence totally enclosed

in the two leaf-sheaths. The important morphological characteristics of each cluster are shown in Table 4. The results were similar to those reported by Siriruga (1992). Based on the morphological characteristics described by Siriruga, 15 *Kaempferia* species from Thailand were divided into two main groups. Eight *Kaempferia* species that included *K. spoliata*, *K. larsenii*, *K. flallax*, *K. filifolia*, *K. elegans*, *K. parviflora*, *K. rotunda*, and *K. angustifolia* were clustered in the first group. These plants have erect leaves together with

elliptic-liner, filliform, lanceolate or orbicular leaf blades. *K. siamensis*, *K. marginata*, *K. galanga*, *K. pulchra*, *K. glauca*, *K. roscoeana*, and *K. laotica* were clustered in the second group which has horizontal leaves.

The results indicated that the genetic relationships through ISSR fingerprints together with morphological characteristics provide a reliable method for identification of plants that have similar morphological characteristics.

Table 4. The important morphological characters of the plants in group I and II.

| Morphological characters | Group I | | Group II | |
|---------------------------|--|--|---|---|
| | Subgroup Ia | Subgroup Ib | Subgroup IIa | Subgroup IIb |
| Height | 6-20 cm | 6-30 cm | 2-5 cm | 2-10 cm |
| Leaf position | Erect | Erect | Horizontal | Horizontal |
| Leaf shape | Elliptic | Elliptic | Elliptic or suborbicular | Elliptic or suborbicular |
| Leaf apex | Acute | Acute | Acute or mucronate | Acute or mucronate |
| Leaf base | Cuneate | Cuneate | Rounded-cuneate | Rounded-cuneate |
| Petiole length | 1-10 cm | Sessile or 1-2 cm | Sessile | Sessile or 3-12 cm |
| Position of inflorescence | Rising between the two innermost leaves or enclose by the two innermost leaf-sheaths | Rising between the two innermost leaves or enclose by the two innermost leaf-sheaths | Enclose by the two innermost leaf-sheaths | Enclose by the two innermost leaf-sheaths |
| Anther crest apex | Entire | Deeply divided | Entire | Deeply divided |

4. Conclusions

The phylogenetic relationships of 11 accessions of *Kaempferia* plants belonging to 10 species from Thailand and Laos were determined by ISSR fingerprints. Forty-five ISSR primers were screened, of which 10 primers produced clear and reproducible bands. According to the ISSR profiles, the *Kaempferia* plants were divided into two major groups. ISSR profiles showed that the ISSR-27 primer generated characteristic bands of Zingiberaceous plants. Therefore, the ISSR-27 primer could be further developed to authenticate Zingiberaceous plants. The ISSR marker is a reliable and cost effective technique in addition to assessing genetic diversity in Zingiberaceous plants. Furthermore, sequence characterized amplified regions could be further developed to quality control of herbal medicines and differentiation of plants in the *Kaempferia* genus. Moreover, this is the first report on ISSR fingerprints of *Kaempferia* plants. The data might be used as additional information for plant conservation and plant breeding for the *Kaempferia* genus.

Acknowledgements

The author would like to acknowledge the Research Institute of Rangsit University for the financial support (Grant no. 39/2557). The author would like to thankful Assoc. Prof. Thatree Phadungcharoen, Department of Pharmacognosy, Faculty of Pharmacy, Rangsit University for the sample collection and valuable information.

References

- Chuakul, W. (2005). Medicinal plants in the Khok Pho District, Pattani Province (Thailand). *Thai Journal of Phytopharmacy*, 12, 23-45.
- Chuakul, W., & Boonpleng, A. (2003). Ethnomedical uses of Thai Zingiberaceous plant (1). *Thai Journal of Phytopharmacy*, 10, 33-39.
- Chuakul, W., & Boonpleng, A. (2004). Survey on Medicinal Plants in Ubon Ratchathani Province (Thailand). *Thai Journal of Phytopharmacy*, 11, 33-54.
- Dogan, B., Duran, A., Seker, M., Cetin, O., & Martin, E. (2015). Study of phylogenetic relationship of Turkish species of *Klasea* (Asteraceae) based on ISSR amplification. *PhytoKeys*, 56, 29-40.
- Dogan, B., Celik, M., Unal, M., Sefali, A., Martin, E., & Kaya, A. (2016). Study of phylogenetic relationship of Turkish species of *Matthiola* (Brassicaceae) based on ISSR amplification. *Turkish Journal of Botany*, 40, 130-136.
- Hao, G., Lee, D., Lee, J. S., & Lee, N. S. (2002). A study of taxonomical relationships among species of Korean *Allium* sect. *Sacculiferum* (Alliaceae) and related species using inter-simple sequence repeat (ISSR) markers. *Botanical Bulletin of Academia Sinica*, 43, 63-68.
- Heubl, G. (2010). New aspects of DNA-based authentication of Chinese medicinal plants by molecular biological techniques. *Planta Medica*, 76, 1963-1974.
- Joshi, S. P., Gupta, V. S., Aggarwal, R. K., Ranjekar, P. K., & Brar, D. S. (2000). Genetic diversity and phylogenetic relationship as revealed by inter-simple sequence repeat (ISSR) polymorphism in the genus *Oryza*. *Theoretical and Applied Genetics*, 100, 1311-1320.
- Kitamura, C., Nagoe, T., Prana, M. S., Agusta, A., Ohashi, K., & Shibuya, H. (2007). Comparison of *Curcuma* sp. in Yakushima with *C. aeruginosa* and *C. zedoaria* in Java by *trnK* gene sequence, RAPD pattern and essential oil component. *Journal of Natural Medicines*, 61, 239-243.

- Maneenoon, K., Khuniad, C., Teuanan, Y., Saedan, N., Prom-In, S., Rukleng, N., . . . Wongwiwat, W. (2015). Ethnomedicinal plants used by traditional healers in Phatthalung Province, Peninsular Thailand. *Journal of Ethnobiology and Ethnomedicine*, 11, 43. doi: 10.1186/s13002-015-0031-5
- Nuamnee, A., Seraypheap, K., Yannawat, S., & Seelanan, T. (2012). Ethnobotany of Hmong at Ban Pang Chang, Pong Subdistrict, Santisuk District, Nan Province. *Thai Journal of Botany*, 4, 177-211.
- Pavlicek, A., Hrda, S., & Flegr, J. (1999). FreeTree-freeeware program for construction of phylogenetic trees on the basis of distance data and bootstrap / jackknife analysis of the tree robustness. Application in the RAPD analysis of genus *Frenkelia*. *Folia Biologica*, 45, 97-99.
- Picheansoonthon, C., & Koonterm, S. (2008). Notes on the genus *Kaempferia* L. (Zingiberaceae) in Thailand. *Journal of Thai Traditional & Alternative Medicine*, 6, 73-93.
- Prevost, A., & Wilkinson, M. J. (1999). A new system of comparing PCR primers applied to ISSR fingerprinting of potato accessions. *Theoretical and Applied Genetics*, 98, 107-112.
- Pooma, R., & Suddee, S. (2014). *Thai plant names (Botanical names - vernacular names) Revised Edition*. Bangkok, Thailand: Royal Forest Department.
- Rout, G. R., Senapati, S. K., & Aparajita, S. (2010). Study of Relationships among Twelve *Phyllanthus* Species with the Use of Molecular Markers. *Czech Journal of Genetics and Plant Breeding*, 46, 135-141.
- Semagn, K., Bjornstad, A., & Ndjondjop, M. N. (2006). An overview of molecular marker methods for plants. *African Journal of Biotechnology*, 5, 2540-2568.
- Siriluck, I., Ratchanok, T., Worakij, H., & Thitamin, K. (2014). Identification of 24 Species of Zingiberaceae in Thailand Using ISSR Technique. *Thai Journal of Agricultural Science*, 47, 1-6.
- Sirirugsa, P. (1992). Taxonomy of the genus *Kaempferia* (Zingiberaceae) in Thailand. *Thai Forest Bulletin*, 19, 1-15.
- Leong-Skornikova, J., & Newman, M. (2015). *Gingers of Cambodia, Laos and Vietnam*. Tanglin, Singapore: Singapore Botanic Garden National Parks Board.
- Son, J., Park, K., Lee, S., Kim, J., & Kim, N. (2012). Species Relationships among *Allium* Species by ISSR Analysis. *Horticulture, Environment, and Biotechnology*, 53, 256-262.
- Suksri, S., Premcharoen, S., Thawatphan, C., & Sangthongprow, S. (2005). Ethnobotany in Bung Khong Long Non-Hunting Area, Northeast Thailand. *Kasetsart Journal: Natural Science*, 39, 519-533.
- Syamkumar, S., & Sasikumar, B. (2007). Molecular marker based genetic diversity analysis of *Curcuma* species from India. *Scientia Horticulturae*, 112, 235-241.
- Tangjitman, K., Wongsawad, C., Kamwong, K., Sukkho, T., & Trisonthi, C. (2015). Ethnomedicinal plants used for digestive system disorders by the Karen of northern Thailand. *Journal of Ethnobiology and Ethnomedicine*, 11, 27. doi:10.1186/s13002-015-0011-9
- Techaprasan, J., Klinbunga, S., Ngamriabsakul, C., & Jentittikul, T. (2010). Genetic variation of *Kaempferia* (Zingiberaceae) in Thailand based on chloroplast DNA (*psbA-trnH* and *petA-psbJ*) sequences. *Genetics and Molecular Research*, 9, 1957-1973.
- Tharachand, C., Immanuel, S. C., & Mythili, M. N. (2012). Molecular markers in characterization of medicinal plants: An overview. *Journal of Plant Biology Research*, 2, 1-12.
- Theanphong, O., Thanakijcharoenpath, W., Ruangrungsi, N., Palanuvej, C., & Rungsihirunrat, K. (2013). Phylogenetic relationship of selected *Kaempferia* plants in Thailand based on RAPD marker. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 4, 1362-1369.
- Vanijajiva, O., Sirirugsa, P., & Suvachittanont, W. (2005). Confirmation of relationships among *Boesenbergia* (Zingiberaceae) and related genera by RAPD. *Biochemical Systematics and Ecology*, 33, 159-170.
- Zou, X., Dai, Z., Ding, C., Zhang, L., Zhou, Y., & Yang, R. (2011). Relationships among six medicinal species of *Curcuma* assessed by RAPD markers. *Journal of Medicinal Plants Research*, 5, 1349-1354.