

Molecular Structure, Electrostatic Potential and HOMO, LUMO Studies of 4-Aminoaniline, 4-Nitroaniline and 4-Isopropylaniline by DFT

Hayat EL Ouafy¹, Mouna Aamor¹, Mustapha Oubenali¹, Mohamed Mbarki¹, Aziz EL Haimouti², Tarik EL Ouafy^{2,*}

 ¹Department of Chemistry and Environment, Faculty of Sciences and Technics, Sultan Moulay Slimane University, Beni Mellal 23000, Morocco
 ²Department of Physics and Chemistry, Polydisciplinary Faculty of Khouribga, Sultan Moulay Slimane University, Beni Mellal 23000, Morocco

Received 26 October 2020; Received in revised form 3 February 2021 Accepted 27 April 2021; Available online 28 March 2022

ABSTRACT

This research used the method of functional density theory (DFT). The B3LYP / 6-311G (d, p) was used to determine the chemical descriptor, the ionization potential (I), the electron affinity (A), the chemical potential (μ), the chemical hardness (η); 3D maps of HOMO and LUMO orbits were used to develop the structure and activity of quantitative relationships. The values of the energy gap showed that p-isopropylaniline is the stable molecule and p-nitroaniline is the reactive molecule. Large basis set-theoretical calculations of the dipole polarizabilities and second hyperpolarizabilities of molecules have been carried out to assess optical properties of atomic contributions to the overall molecular response tensors. The values of the dipole moment and the first-order hyperpolarizability have the same tendency because they are related to molecular symmetry. The DFT method has been used of which is to compare the angles and lengths of molecular bonds with the experimental results. This research used the electrostatic molecular potential (MEP) and electrostatic contour to understand the regions of reactivity of the three molecules. The negative molecular electrostatic potential sites intended for electrophilic attack for p-aminoaniline, p-nitroaniline and p-isopropylaniline are (C1, C2, C3, C4, C5, C6, N7, N8), (O9, O10) and (C5, C6), respectively.

Keywords: 4-Aminoaniline; 4-Nitroaniline; 4-Isopropylaniline; DFT; Chemical descriptor; NLO; Electrostatic molecular potential

1. Introduction

The density functional theory (DFT) plays an important role in studies on freezing and melting transitions [1-2]. The liquid-crystal interface [3], nucleation [4], glass transition [5] and quasi-crystals [6] are also treated within the framework of the DFT.

Aromatic amines play an important role in the chemical and biological industries. Aniline and its derivatives are utilized in the production of pesticides,

dyes, and antioxidants; some of the psubstituted derivatives of aniline are local anesthetics. Aniline, also known as aminobenzene, phenylamine, or benzene amine, is an aromatic organic compound with the chemical formula C_6H_5 -NH₂ [7]. 4-Amino-aniline (Fig. 1A) represents a very interesting class of products possessing a p and r electron system and its derivatives are used in the synthesis of dyes. Its ability to donate electrons is explored in combination with an electron acceptor. It is important for us to determine the properties of its best nature as an electron donor [8]. Aromatic amines are characterized by importance in science and materials as well as in the chemical and pharmaceutical fields. Due to their high toxicity and wide uses, halogenated anilines have obtained the highest concentration of all aromatic amines. Para-nitroaniline (Fig. 1B), or 1amino-4-nitrobenzene, is chemical а compound with the formula $C_6H_6N_2O_2$. It is an almost odorless, yellowish-orange, crystalline powder, combustible and very soluble in water, where it is very toxic to aquatic organisms [9]. Amino and methyl groups are usually called electron giving substituents to aromatic ring systems. The CH₃ group reacts with neighboring p systems via hyperconjugation, while the NH₂ group reacts with the p electrons in a ring. Both reactions involve delocalization

of electrons. Toluidine (Fig. 1C), or aminotoluene is an aromatic compound of formula C7H9N, consisting of a benzene ring substituted by an amine group (aniline) and a methyl group (toluene). Like all disubstituted benzenes, it exists in the form of three structural isomers, the ortho, meta and para compounds, depending on the relative position of the two substituents on the ring [10]. Charge transfer interactions form between electron donor compounds and electron acceptor compounds and result in resonance with charge transfer from the donor molecule to the acceptor molecule. Complex formation is based on the interaction of the donor's highest occupied molecular orbital with the lowest unoccupied acceptor molecular orbital [11].

This research is interested in studying the structural properties of p-aminoaniline, p-nitroaniline and 4-isopropylaniline. The DFT method was used to calculate the global descriptors (the energies and densities of the limit molecular orbitals HOMO and LUMO, the electronic chemical potential, electronegativity, chemical hardness, overall softness, overall electrophilic index, global nucleophilic index, properties nonlinear magnetic optics) to compare the stability and reactivity of these molecules and determine their regions of reactivity by molecular electrostatic potential.



Fig. 1. Optimized molecular structure of (A) p-aminoaniline, (B) p-nitroaniline, (C) p-isopropylaniline.

2. Computational Method

All calculations have been carried out with the Gaussian 09 program [12] and visualized with the Gaussview software [13] on an i7 PC. The optimization was performed by density functional theory (DFT) [14] with the B3LYP functional [15-16] and the standard base 6-311G (d, p).

The DFT method was used to include popular local methods and gradient correction [17]. The following quantum of the chemical indices have been taken into account: the E_{HUMO} , E_{LUMO} , the energy band gap $\Delta E = E_{HUMO} - E_{LUMO}$, the quantum descriptors, the electronic affinity, the ionization potential and the atomic partial charges were calculated for the two structures from the potential electrostatic surface (ESP) according to the same theory of levels [18].

3. Results and Discussion 3.1 Ouantum chemical calculation

The main orbitals involved in chemical stability are the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO). The HOMO represents the capacity to yield an electron while the LUMO represents the capacity to gain an electron. The HOMO and LUMO energies are calculated by the B3LYP/6-311G (d, p) method. Electronic absorption corresponding to the passage of the ground to the first excited state described by an electronic excitation from the HOMO orbital to the LUMO orbital [19] with the optimized structure of molecules is illustrated in Figs. 2-3 shows the Mulliken charge distribution for the three molecules.

номо



Fig. 2. HOMO and LUMO of (A) p-aminoaniline, (B) p-nitroaniline, (C) p-isopropylaniline.



Fig. 3. Mulliken charges of (A) p-aminoaniline, (B) p-nitroaniline, (C) p-isopropylaniline by the B3LYP method.

Many applications are available on the utilization of the HOMO and LUMO gap as a quantum descriptor in correlates in different biochemical and chemical systems [20].

The following quantum descriptors have been calculated from the optimized structure obtained: Ionization potential:

$$I = -E_{HUMO}, \qquad (3.1)$$

Electronic affinity:

$$A = -E_{LUMO}, \qquad (3.2)$$

Absolute electronegativity:

$$\chi = \frac{I+A}{2},\tag{3.3}$$

Overall hardness:

$$\eta = I - A, \tag{3.4}$$

Overall softness:

$$\sigma = \frac{1}{\eta} = \frac{1}{E_{LUMO} - E_{HOMO}},\qquad(3.5)$$

Electronic chemical potential:

$$\mu = -\frac{\left(I+A\right)}{2},\tag{3.6}$$

Maximum charge transfer:

$$\Delta N_{\rm max} = \frac{\mu}{n}, \qquad (3.7)$$

Overall electrophilicity:

$$\omega = \frac{\mu^2}{2\eta},\tag{3.8}$$

Overall nucleophilia N:

$$N = E_{HUMO} - E_{HOMO(TCE)}, \qquad (3.9)$$

with $E_{HOMO(TCE)} = -9.3686$ eV calculated by DFT/B3LYP 6-311G (d, p).

In order to highlight the electrophilic/nucleophilic character of the molecules, we calculated the ionization potential I, the electronic affinity A, the electronic chemical potential μ , the absolute electronegativity χ , the overall hardness η , the overall softness σ , the overall electrophilicity index ω , the index of overall nucleophilic N and the maximum charge transfer ΔN_{max} which are shown in Table 1.

Table 1. Quantum theoretica	l parameters of (A)	p-aminoaniline,	(B) p-nitroaniline,	(C) p-iso-
propylaniline, calculated by H	B3LYP / 6-311G (d	, p).		

Parameters	Elumo(eV) Ehomo	(eV) Δ	E (eV)	I (eV)	A (eV) µ (eV)
p-Aminoaniline	0.230	6 -4.3	713	4.6019	4.3713	-0.230	6 -2.0703
Parameters	χ (eV)	η (eV)	σ (eV ⁻¹)	ω (eV)		N (eV)	$\Delta N_{max}(eV)$
p-Aminoaniline	2.0703	4.6019	0.2173	0.6456		4.9973	0.4498
Parameters	E _{LUMO} (eV)	E _{HOMO} (eV)	$\Delta E (eV)$	I (eV)		A (eV)	μ (eV)
p-Isopropylaniline	0.0682	-5.2286	5.2968	5.2286		-0.0682	-2.5802
Parameters	χ (eV)	η (eV)	σ (eV ⁻¹)	ω (eV)	1	N (eV)	ΔN_{max} (eV)
p-Isopropylaniline	2.5802	5.2968	0.1887	0.6284	4	4.14	0.4871
Parameters	E _{LUMO} (eV)	E _{HOMO} (eV)) ΔE (e	eV) I (e	V)	A (eV)	μ (eV)
p-Nitroaniline	-2.5714	-6.4621	3.890	6.4	621	2.5714	-4.5167
Parameters	χ (eV)	η (eV)	σ (eV ⁻¹)	ω (eV)	Ν	(eV)	$\Delta N_{max}(eV)$
p-Nitroaniline	4.5167	5.7313	0.1744	1.7797	2.	9065	0.7880

The calculated energy gap values are 3.8907 eV, 5.2968 eV and 4.6019 eV for pnitroaniline, p-isopropylaniline and p-aminoaniline, respectively. This explains why p-isopropylaniline is more stable than p-nitroaniline and p-aminoaniline.

The overall electrophilicity index values are 0.6456 eV, 0.6284 eV and 1.7797 eV for p-aminoaniline, p-isopropy-laniline and p-nitroaniline, respectively. As a result, the reactivity of p-nitroaniline to a nucleophilic attack is greater than the other two molecules.

The overall nucleophilicity index values are 4.14 eV, 2.9065 eV and 4.9973 eV, for p-isopropylaniline, p-nitroaniline and p-aminoaniline. Therefore, the reactivity of p-aminoaniline to electrophilic attack is greater than that of p-nitroaniline and pisopropylaniline.

The charge that p-nitroaniline can acquire from its environment ($\Delta N_{\text{max}} = 0.7880 \text{ eV}$) is greater than that of other molecules.

3.2 Non-linear optical properties

Intermolecular interactions such as drug-vitamin are widely understood by dipole moment and the energetic terms of hyperpolarization of the first and second-order. The dipole moment (μ), the polarizability (α), the first hyperpolarizability (β) and the second hyperpolarizability (γ) are calculated using DFT on the basis B3LYP 6-311G approach (d, p). The complete equations to calculate the amplitude of the total static dipole moment (μ), the polarizability (α), the first hyperpolarizability (β) and the second hyperpolarizability (γ), by the components *x*, *y*, *z* of 09W the Gaussian output is as follows [21]:

$$\mu = \left(\mu_x^2 + \mu_y^2 + \mu_z^2\right)^{\frac{1}{2}}, \qquad (3.10)$$

$$\alpha = \frac{\alpha_{xx} + \alpha_{yy} + \alpha_{zz}}{3}, \qquad (3.11)$$

$$\beta = \left(\beta_x^2 + \beta_y^2 + \beta_z^2\right)^{\frac{1}{2}},$$
(3.12)

$$\beta_x = \beta_{xxx} + \beta_{xyy} + \beta_{xzz}, \qquad (3.13)$$

$$\beta_{y} = \beta_{yyy} + \beta_{xxy} + \beta_{yzz}, \qquad (3.14)$$

$$\beta_z = \beta_{zzz} + \beta_{xxz} + \beta_{yyz},$$
(3.15)

$$\langle \gamma \rangle = \frac{1}{5} \Big(\gamma_{xxxx} + \gamma_{yyyy} + \gamma_{zzzz} + 2 \Big[\gamma_{xxyy} + \gamma_{yyzz} + \gamma_{xxzz} \Big] \Big).$$
(3.16)

The results for the non-linear optical properties are tabulated in Table 2.

The dipole moment of the molecules is again calculated using DFT and method B3LYP with basic set 6-311G (d, p). The dipole moment reflects the distribution of molecular charges and is given as a threedimensional vector. Consequently, it can be utilized as a descriptor to represent the charge movement through the molecule as a function of the negative and positive charge centers. Dipole moments are necessarily determined for neutral molecules. For charged molecules, values depend on the orientation and the choice of the origin of the molecular.

The results showed that the determined value of the dipole moment of p-nitroaniline (10.0817 D) is greater than those of p-isopropylaniline (2.4864 D) and paminoaniline (0.0003 D).

The results showed that the dipolar polarizability value of p-aminoaniline (-42.3552 D) was higher compared to the values of the polarizability tensors of p-nitroaniline and p-isopropylaniline.

The transitions of p-nitroaniline 125.9916 D and -0.0007 D are observed

along the x and y axes and -0.0008 D along with a negative direction, while the transitions of p-aminoaniline -0.0008 D and -0.0009 D are found along the y and z axes with a negative direction and the 0.0008 D transition is found along with a positive direction. The p-isopropylaniline transitions -0.8625 D and -1.0064 D are found along the y and z axes with a negative direction and 78.4461 D along the x-axis with a positive direction.

Dipole and first order hyperpolarizability have the same tendency because they are related to molecular symmetry.

The high and low value of β is calculated to be 135.1874 D and 0.00475 in pnitroaniline and aminoaniline, respectively. Therefore, electron delocalization and charge transfer increase in p-nitroaniline, resulting in a greater β response of p-nitroaniline compared to p-isopropylaniline and p-aminoaniline.

The values of the second-order hyperpolarizability γ , as well as their contributing tensors, are also observed to be greater in p-aminoaniline then p-nitroaniline and p-isopropylaniline.

Table 2. Electric dipole moments (Debye) by the DFT method of (A) p-aminoaniline, (B) pnitroaniline, (C) p-isopropylaniline calculated by B3LYP / 6-311G (d, p).

	Parameters	Α	В	С
	μ_x	0.0000	10.0817	2.4794
Dipole moment (Debye)	μ_{y}	-0.0002	-0.0002	-0.1621
	μ_{z}	0.0002	0.0001	0.0932
	μ	0.0003	10.0817	2.4864
D _1	α_{xx}	-30.5966	-56.3980	-48.7801
Polarizability (Debye)	α_{yy}	-42.1883	-53.5752	-58.3983
	α_{zz}	-54.2808	-60.0852	-67.1977
	α	-42.3552	-56.6861	-58.1253
	$\beta_{_{\rm XXX}}$	-0.0008	125.9916	78.4461
First Hyperpolarizability (Debye)	$\beta_{_{xxy}}$	0.0002	19.8964	13.8498
	$\beta_{_{xzz}}$	0.0004	-11.1622	-0.2943
	$\beta_{_{VVV}}$	-0.0009	-0.0007	-0.8625
	$\beta_{_{xxy}}$	-0.0022	-0.0001	-0.5293
	$\beta_{_{VZZ}}$	0.0000	0.0002	-0.4765
	$\beta_{_{777}}$	0.0008	-0.0008	-1.0064

H. EL Ouafy et al.	Science &	Technology 2	1sia Vol.27 No.	1 January - March 2022	
			0.001=	11.1.000	0

	$\beta_{_{xxz}}$	0.0017	-11.1622	-0.2943
	$eta_{_{yyz}}$	0.0011	0.0003	1.2485
	β	0.00475	135.1878	92.0205
	γ_{xxxx}	-631.3569	-1246.4404	-1472.2424
Second	γ_{yyyy}	-308.6713	-370.0793	-392.4747
Hyperpolarizability (Debve)	γ_{zzzz}	-58.9777	-59.6126	-220.9768
(~ , -)	γ_{xxyy}	-145.4924	-285.8703	-326.6312
	$\gamma_{_{VVZZ}}$	-72.3390	-83.5331	-108.2036
	γ _{xx77}	-175.3008	-253.8500	-352.9236
	γ	-357.0540	-584.5277	-732.2421

3.3 Thermodynamic parameters

This research determined for each molecule the enthalpy ΔH , the entropy ΔS , and the free energy ΔG (Table 3). The enthalpy ΔH is defined by the sum of the internal energy of a system and the product of its pressure by its volume. Entropy ΔS can be characterized by the degree of disorder or unpredictability of the information contained in a system. The ΔG is the energy rel-

ative to different reference states. The values of Δ H and Δ S were taken directly from the Notepad file provided by the Gaussian software and Δ G was calculated by Δ G = Δ H-T Δ S with T = 298K. This Δ G amount is relative to free electrons and free nuclei. We found that the Δ G values of p-nitroaniline, p-aminoaniline and p-isopropylaniline are -1291.5 × 10³ kj/mol, -899.9 × 10³ kj/mol and -1064.3 × 10³ kj/mol, respectively.

Table 3. Thermodynamic parameters of (A) p-aminoaniline, (B) p-nitroaniline, (C) p-isopropvlaniline

Molecules	∆H (Kj/mol)	ΔS (Kj/mol/K)	$\Delta \mathbf{G}$ (Kj/mol)
Α	-900×10^{3}	0.36024	-899.9×10^{3}
В	-1291.6×10^{3}	0.37278	-1291.5×10^{3}
С	-1064.2×10^{3}	0.41045	-1064.3×10^{3}

3.4 Bond length and bond angle properties

This research will deepen our knowledge of selective bond lengths (Å) and angles (θ°). The DFT level has been used with the B3LYP/6-311G (d, p) method, the utility of which is to describe the molecular physical and chemical properties [18].

Tables 4-5 were determined the comparison of the optimized bond length and the angle between experimental and calculated atomic numbered positions as shown in Fig. 1. The optimized structures using a periodic DFT calculation correspond exactly to an experimental result. The study of the geometric product, therefore, showed an excellent agreement between theoretical and experimental results.

Fable 4. Bond lengths (Å) of (A)	p-aminoaniline, (B)	p-nitroaniline,	(C)	p-isopropylaniline) .
-------------------------	-----------	-------------------	------------	-----------------	-----	--------------------	------------

Bond length (Å)	Α	В	С	Exp [7,22]
C1-C2	1.3948	1.3948	1.3948	1.408
C2-C3	1.3949	1.3949	1.3949	1.415
C3-C4	1.3948	1.3948	1.3948	1.390
C4-C5	1.3948	1.3948	1.3948	1.395
C5-C6	1.3949	1.3949	1.3949	1.395
C6-N7	1.266	1.266	1.266	1.370
C3-N8	1.266			1.247

H. EL Ouafy et al. | Science & Technology Asia | Vol.27 No.1 January - March 2022

C3-O8	 1.248		1.246
N8-O9	 1.31		
N8-O10	 1.31		
C3-C8	 	1.497	1.460
C8-C9	 	1.523	
C8-C10	 	1.523	1.737

Tuble 5. Thisles (6) of (11) p anniounnine, (b) p introunnine, (c) p isopropyiunnine.	Table 5. Angles	(θ^{o})) of (A)	p-aminoai	niline, (B)	p-nitroaniline,	(C)	p-isopropylaniline.
---	-----------------	----------------	----------	-----------	-------------	-----------------	-----	---------------------

Angles (θ°)	Α	В	С	Exp [7]
C1-C2-C3	119.9969	119.9969	119.9969	118.8
C2-C3-C4	120.0002	120.0002	120.0002	120.9
C3-C4-C5	120.0033	120.0033	120.0033	120.0
C5-C6-N7	119.9991	119.9991	119.9991	119.5
C2-C3-N8	120.0034	119.9973		120.4
O9N8-O10		120.0008		120.8
C9C8-C10			109.5028	120.1

3.5 Electrostatic Molecular Potential of molecules

The electrostatic molecular potential gives detailed information for the study of the biological activity and chemical reactivity of a molecule. The distribution and values of electrostatic potential determine the sites of electrophilic or nucleophilic attack in a chemical reaction [22-27]. The MEP is mainly used in the form of the reactivity map showing the most likely regions for the electrophile attack of point reagents loaded on organic compounds. It is important in the molecular modeling studies. Contour MEP, the map offers a simple tool to predict how different geometries can interact. The total electronic density and the MEP surface of the molecules studied are constructed by B3LYP / 6-311 method using

basis set G (d, p). Electrostatic potential maps and contour electrostatic potential of p-aminoaniline, p-nitroaniline and isopropylaniline are shown in Figs. 4 and 5. The color scheme for the MEP surface is red, rich in electrons, partially negative charge; blue, electron-deficient, partially positive charge; light blue, slightly electron-deficient region; yellow region, slightly rich in electrons; green, neutral, respectively. Blue represents electropositive and red the electronegative regions, respectively [28].

The negative molecular electrostatic potential sites intended for electrophilic attack for p-aminoaniline, p-nitroaniline, and p-isopropylaniline are (C1, C2, C3, C4, C5, C6, N7, N8), (O9, O10) and (C5, C6), respectively.



Fig. 4. Electrostatic potential maps around the molecule of p-aminoaniline, p-nitroaniline and p-isopropylaniline.

Fig. 5. Contour electrostatic potential around the molecule of p-aminoaniline, p-nitroaniline and p-iso-propylaniline.

4. Conclusions

The electronic properties of the three molecules have been calculated. HOMO and LUMO energy deviations describe the possible charge, transfer interactions taking place in the molecule, and the determination of theoretical molecular structures, electrostatic potential and electrostatic potential contour maps around molecules by B3LYP / 6-311G (d, p) using the DFT method. Molecular orbitals, the MEP surface can lead to an understanding of the properties and activity of molecules. The linear polarizability (α), first hyperpolarizability (β) and second hyperpolarizability (γ) values of the studied molecule were calculated. DFT / B3LYP / 6-311G (d, p) has proven to be a good method of optimizing molecular geometries. The molecular electrostatic potential determined the regions rich in electrons destined for electrophilic attacks.

References

- [1] Scuseria GE, Janssen CL, Schaefer III, HF. An efficient reformulation of single and double excited equations coupled to a closed-shell cluster (CCSD). J Chem Phys 1988;89:7382-7.
- [2] Miehlich B, Savin A, Stoll H, Preuss H. Results obtained with the correlation energy density functionals of Becke and

Lee, Yang and Parr. Chem Phys Lett 1989;157:200-6.

- [3] Lipari NO, Duke CB. The electronic structure of dialkylbenzenes. J Chem Phys 1975;63:1748-57.
- [4] Purvis III GD, Bartlett RJ. A complete coupled single and double cluster model: the inclusion of disconnected triples. J Chem Phys 1982;76:1910-18.
- [5] Norman ROC. Taylor R. Electrophilic substitution in benzenoid compounds. Elsevier Publishing Company 1963;3:3562-70.
- [6] McLean AD. Chandler GS. Sets of contracted Gaussian bases for molecular calculations. I. Second row atoms, Z = 11-18. J Chem Phys 1980;72:5639-48.
- [7] Kavitha E, Sundaraganesan N, Sebastian S. Molecular structure, vibrational spectroscopic and HOMO, LUMO studies of 4-nitroaniline by density functional method. J pur appl phys 2010;48:20-30.
- [8] Lakkadi A, Baindla N, Tigulla P. Synthesis, spectroscopic and computational studies of charge-transfer complexation between 4-aminoaniline and 2, 3-dichloro-5, 6-dicyano-1, 4-benzoquinone. J Sol Chem 2017;46:2171-90.

- [9] Ribeiro da Silva MA, Ferreira AI, Gomes JR. Experimental and computational study on the thermochemistry of bromoanilines. Bulletin of the Chemi Soci Japan 2017;79:1852-9.
- [10] Arivazhagan M, Rexalin DA. FT-IR, FT-Raman, NMR studies and ab initio-HF, DFT-B3LYP vibrational analysis of 4-chloro-2-fluoroaniline. Spectro Acta Part A: Molec Biomole Spectro 2012;96:668-76.
- [11] Zeroual A, Ríos-Gutiérrez M, Amiri O, Domingo LR. A molecular electron density theory study of the mechanism, chemo-and stereoselectivity of the epoxidation reaction of R-carvone with peracetic acid. RSC Advances. 2019;9:28500-9.
- [12] Gaussian09 R A, Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Mennucci B, Petersson GA. gaussian. Inc., Wallingford CT. 2009;121:150-66.
- [13] Frisch A, Nielson AB, Holder AJ. Gaussview user manual. Gaussian Inc., Pittsburgh, PA, 2000;556.
- [14] Hohenberg P, Kohn W. Inhomogeneous electron gas. Phys Rev. 1964;136:B864-B871.
- [15] Becke AD. A new mixing of Hartree– Fock and local density-functional theories. J Chem Phys. 1993;98:1372-7.
- [16] Lee C, Yang W, Parr RG. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. Phys Rev B. 1988;37:785-7.
- [17] Suresh S, Gunasekaran S, Srinivasan S. Spectroscopic (FT-IR, FT-Raman, NMR and UV-Visible) and quantum chemical studies of molecular geometry, Frontier molecular orbital, NLO, NBO and thermodynamic properties of salicylic acid. Spectrochim Acta - Part A Mol Biomol Spectrosc 2014;132:130-41.

- [18] Dobado JA, Molina J. Adenine-Hydrogen Peroxide System: DFT and MP2 Investigation. J Phys Chem A 1999;103:4755-61.
- [19] Kavitha E, Sundaraganesan N, Sebastian S, Kurt M. Molecular structure, anharmonic vibrational frequencies and NBO analysis of naphthalene acetic acid by density functional theory calculations. Spectrochim Acta - Part A Mol Biomol Spectrosc 2010;77:612-19.
- [20] Sebastian S, Sylvestre S, Jayabharathi J, Ayyapand S, Amalanathan M, Oudayakumar K. Study on conformational stability, molecular structure, vibrational spectra, NBO, TD-DFT, HOMO and LUMO analysis of 3,5-dinitrosalicylic acid by DFT techniques. Spectrochim. Acta - Part A Mol Biomol Spectrosc 2015;136:1107-18.
- [21] Liu L, Gao H. Molecular structure and vibrational spectra of ibuprofen using density function theory calculations. Spectrochim. Acta - Part A Mol Biomol Spectrosc 2012;89:201-9.
- [22] Mohammad AM, Tarun KP, Ashraful A, Al-Amin-Al-Azadul I, Subrata P, Chanmiya S. DFT studies on vibrational and electronic spectra, HOMO–LUMO, MEP, HOMA, NBO and molecular docking analysis of benzyl-3-N-(2,4,5trimethoxyphenylmethylene)hydrazinecarbodithioate. J Mol Stru 2020;1220:128715-49.
- [23] Mps TF. Representation of molecular electrosatic potentials by topological feature maps. J Am Chem SOC 1994;116:4608-20.
- [24] EL Ouafy H, Aamor M, Oubenali M, Mbarki M, Gamouh A, EL Haimouti A, EL Ouafy T. 4-Bromo-N,N-dimethylaniline, 4-fluoro-N,N-dimethylaniline, 4-methyl-N,N-dimethylaniline: Densityfunctional theory study. Sci Eng Health Studies 2021;15: 1-8.
- [25] EL Ouafy H, EL Ouafy T, Oubenali M, Mbarki M, Echajia M. Analysis of the molecular electrostatic potential of the chemical reactivity of p-nitrophenol, paminophenol and p-methylphenol by the

quantum method. J Envir Eng Sci 2020;6:363-73.

- [26] EL Ouafy H, EL Ouafy T, Oubenali M, Mbarki M, EL Haimouti A, Echajia M. Application of the quantum chemical descriptor to the quantitative structure-activity of the ascorbic acid molecule: DFT and MP2 survey. Arab J Chem Envir Res 2021;8:58-69.
- [27] EL Ouafy H, EL Ouafy T, Oubenali M, Mbarki M, EL Haimouti A, Echajia M. Analysis of the Chemical Reactivity of

Limonene by the Functional Density Theory Method Using Global Descriptors. J Chem Healt Ris 2021;11: 10.22034/jchr.2021.1910282.1189.

[28] Hathwar VR, Pal R, Guru Row TN. Charge density analysis of crystals of nicotinamide with salicylic acid and oxalic acid: An insight into the salt to cocrystal continuum. Cryst Growth Des 2010;10:3306-10.