

Molecular Characterization of *Wenyonella philiplevinei* from Ducks in Hunan Province, China

Hui-Lan Wu ^{1*} Peng-Hui Hu ² Jin Zhang ³

Abstract

Wenyonella philiplevinei is one of the important pathogens of duck coccidiosis. In the present study, a portion of the 18S rDNA was amplified from *W. philiplevinei* by Polymerase chain reaction (PCR), cloned and then sequenced. The sequences of 18S rDNA of all samples were 422 bp in size. A + T content of the 18S rDNA sequences was 58-59%. Sequence comparison revealed that the similarity in 18S rDNA sequences between Hunan isolates and *W. philiplevinei* available (Guangdong isolate) was more than 98%. Intra-specific sequence variations within each of the Hunan isolates were 0-1.7%. However, the inter-specific sequence differences between *W. philiplevinei* and coccidian available in GenBank were more than 7%. Phylogenetic analysis using neighbour joining (NJ), maximum likelihood (ML) and maximum parsimony (MP) methods indicated that the genus *Wenyonella* was more closely related to *Eimeria* + *Cyclospora* than to *Isospora*. These novel data provide useful genetic marker for the differentiation of *W. philiplevinei* or other closely related coccidians.

Keywords: 18S rDNA, duck, phylogenetic analysis, *Wenyonella philiplevinei*

¹Hunan First Normal University, Changsha 410205, China

²Center for Quality Quarantine of Livestock, Poultry and Aquaculture Products, Changsha 410128, China

³Supervision of Animal Epidemic Prevention Station of Hunan, Changsha, 410013, China

*Correspondence author E-mail: wuhuilan2013@163.com

บทคัดย่อ

ลักษณะทางชีวโมเลกุลของ *Wenyonella philiplevinei* จากเป็ดในมณฑลหูหนานประเทศจีน

Hui-Lan Wu^{1*} Peng-Hui Hu² Jin Zhang³

Wenyonella philiplevinei เป็นหนึ่งในเชื้อที่สำคัญที่ก่อโรคบิดในเป็ด การศึกษาครั้งนี้ ส่วนหนึ่งของ 18S rDNA ของ *W. philiplevinei* ถูกเพิ่มปริมาณโดยปฏิกิริยาลูกโซ่โพลีเมอร์เรส (PCR) จากนั้นทำการโคลนและวิเคราะห์ลำดับเบส ลำดับของ 18S rDNA ของทุกตัวอย่างมีขนาด 422 bp ปริมาณ A และ T ของ 18S rDNA มีค่าร้อยละ 58-59 การเปรียบเทียบลำดับเบสแสดงให้เห็นว่ามีความคล้ายคลึงกันในระดับ 18S rDNA ระหว่างเชื้อที่แยกได้จากห่าน และ *W. philiplevinei* (จาก Guangdong) มากกว่าร้อยละ 98 ความแตกต่างที่จำเพาะในแต่ละสายพันธุ์ของเชื้อที่แยกจากห่านคิดเป็นร้อยละ 0-1.7 อย่างไรก็ตามความแตกต่างที่จำเพาะระหว่าง *W. philiplevinei* และ coccidian ที่มีข้อมูลใน GenBank มีมากกว่าร้อยละ 7 การวิเคราะห์สายวิวัฒนาการ ด้วยวิธี Neighbour Joining (NJ), Maximum Likelihood (ML) และ Maximum Parsimony (MP) ชี้ให้เห็นว่า สายพันธุ์ *Wenyonella* มีความสัมพันธ์อย่างใกล้ชิดกับ *Eimeria* และ *Cyclospora* มากกว่า *Isospora* ข้อมูลใหม่เหล่านี้มีประโยชน์ในแง่ของการใช้เป็นเครื่องหมายทางพันธุกรรมสำหรับหาความแตกต่างของ *W. philiplevinei* หรือเชื้อบิดอื่นๆ ที่เกี่ยวข้อง

คำสำคัญ: 18S rDNA เป็ด phylogenetic analysis *Wenyonella philiplevinei*

¹ Hunan First Normal University, Changsha 410205, China

² Center for Quality Quarantine of Livestock, Poultry and Aquaculture Products, Changsha 410128, China

³ Supervision of Animal Epidemic Prevention Station of Hunan, Changsha, 410013, China

*ผู้รับผิดชอบบทความ E-mail: wuhuilan2013@163.com

Introduction

Coccidiosis in ducks can be caused by members of the genera *Eimeria*, *Isospora*, *Tyzzeria* and *Wenyonella*, and seriously impairs growth and feed utilization of infected ducks resulting in production losses (Edgar, 1987). Coccidian is highly prevalent in ducks and *W. philiplevinei* is considered as predominant species in China (Yin et al., 1983). Previous investigations (Huang et al., 2004) showed that the *W. philiplevinei* is highly prevalent (30-80%) in many provinces or regions of China, including Henan, Xinjiang, Shanghai, Fujian, Yunan, Jiangxi, Beijing, Jiangsu and Guangdong. Although *W. philiplevinei* is considered a significant pathogen in ducks in China, until now, there is only one report of molecular characterization of *W. philiplevinei* (Chen et al., 2010). Furthermore, *W. philiplevinei* is also well recognized as the significant protozoan pathogens affecting poultry industry throughout the world (Skirnisson, 1997). However, little is known about the prevalence of *W. philiplevinei* in duck in other countries and regions. *W. philiplevinei* can cause serious damage to the digestive tract of the host, resulting in decreased bodyweight gain and diarrhea, and can even lead to death (Yin et al., 1983). Due to its high mortality, reduced production and control costs (Tewari and Maharana, 2011), the prevention of duck coccidiosis poses a challenge for the poultry industry development. Moreover, to our knowledge, no information is available about phylogenetic relationships of *W. philiplevinei*. The objectives of the present study were to characterize 18S rDNA sequences of the *W. philiplevinei* in ducks from China.

Moreover, based on the 18S rDNA sequences, the phylogenetic relationships in *W. philiplevinei* were also reconstructed.

Materials and Methods

Parasites and isolation of genomic DNA: Oocysts of *W. philiplevinei* were obtained from feces of ducks in Linwu county, Hunan Province, China. The oocysts of *W. philiplevinei* were washed in physiological saline, identified primarily based on morphological characters and predilection sites to species (Zhang et al., 1999), and then were stored in 2.5% potassium dichromate at 4°C. Purified oocysts of the *W. philiplevinei* were washed in phosphate buffered saline (PBS), disrupted in glass beads, and the total genomic DNA was extracted from individual samples by sodium dodecyl sulphate/proteinase K treatment, column-purified (WizardTM DNA Clean-Up, Promega) and eluted into 50 µl distilled water according to the manufacturer's recommendations.

Enzymatic amplification: A portion of the 18S rDNA was amplified with primers Q1 and Q2 (Chen et al., 2010). These primers were synthesised on a Biosearch Model 8700 DNA synthesizer (Shanghai, China). PCR reactions (25 µl) were performed in 2 mM of MgCl₂, 2.5 µM of each primer, 2.5 µl 10×rTaq buffer, 0.2 mM of each dNTPs, 1.25 U of rTaq DNA polymerase (Takara, Japan), and 2 µl of DNA sample in a thermocycler (BioRad, USA) under the following conditions: after an initial denaturation at 94°C for 5 min, then 94°C for 30 sec (denaturation); 48°C for 30 sec (annealing); 72°C for 1 min (extension) for 35 cycles, followed by a final extension at 72°C for 10

min. Five microlitres of each amplicon was examined by 0.8% (w/v) agarose gel electrophoresis to validate amplification efficiency. PCR products were sent to Sangon Company (Shanghai, China) for sequencing for both directions.

Sequences analysis and phylogenetic reconstruction: Sequences of the 18S rDNA were separately aligned using Clustal X 1.83 computer program (Thompson et al., 1997). The sequences of 18S rDNA available in this study were used for phylogenetic analyses. Three methods, namely neighbour joining (NJ), maximum likelihood (ML) and maximum parsimony (MP), were used for phylogenetic re-constructions. NJ analysis was carried out using Dayhoff matrix model implemented by MEGA 4.0 (Tamura et al., 2007). ML analysis was performed using PUZZLE 4.1 under default setting (Strimmer and Haeseler, 1996) and Standard unweighted MP was performed using package Phylip 3.67 (Felsenstein., 1995). Consensus tree was obtained after bootstrap analysis, with 100 for ML tree, and 1000 replications for NJ and MP trees, with values above 50% reported. To study the genetic relationships among Eimeriidae coccidians, other Eimeriidae coccidians were considered in the present study (*Eimeria adeneodei* AF324212; *Eimeria meleagridis* KC305200; *Eimeria trichosuri* FJ829323; *Eimeria scholtysecki* AF324216; *Isospora suis* ISU97523; *Isospora felis* ISRRGE; *Cyclospora colobi* AF111186; *Cyclospora papionis* AF111187), with *Cryptosporidium parvum* (GenBank accession no: AF108864) as the outgroup. Phylograms were drawn using the Tree View v. 1.65 program (Page, 1996).

Results and Discussion

In the present study, specimens of *W. philiplevinei* infecting ducks from China were characterized on the bases of sequences of 18S rDNA regions. Many previous studies have indicated that 18S rDNA sequences can provide useful genetic markers for accurate identification and differentiation of coccidian species (Wünschmann et al., 2010; Ogedengbe et al., 2011; Carlson-Bremer et al., 2012).

In the present study, genomic DNA was extracted from 10 individual *W. philiplevinei* in Hunan province, China. Sequences of the 18S rDNA (~ 430 bp) were amplified individually and subjected to agarose gel electrophoresis (Fig 1). The results indicated that no size mutation was detected on agarose gels among any of the amplicons examined, and no products were also amplified from host DNA.. The sequences of 18S rDNA were 424 bp in length. The A + T content of the 18S rDNA sequences was 58-59%. Sequence comparison revealed that the similarity in 18S rDNA sequences between Hunan isolates and the *W. philiplevinei* available (Guangdong isolate) were more than 98%. The intra-specific sequence variations within each of the Hunan isolates were 0-1.7% (Fig 2). However, the inter-specific sequence differences between *W. philiplevinei* and the coccidian available in GenBank were more than 7%. This result was consistent with that of a previous study (Chen et al., 2010).




Figure 1 PCR-amplified results of 18S rDNA from *Wenyonella philiplevinei* samples by agarose gel electrophoresis. M: DL2000 DNA marker; 1-10: WP1-W10; 11: Negative control.

Chen	TTTCGTGCGTGGTATCCGCTCCGCTTTACAGTGGATACCTGGTTCGCCCTTGGCAATTTTCGGGATCTTTCCCACTTAATTTGTTGGGTTTGATTTCGGAACATTACTTTGAGAAAAATAGAGTGTTCGAAGCGCTTGTGCG	150
WP1		150
WP8		150
WP10		150
WP4		150
WP3	T.....T.....	150
WP6	T.....T.....	150
WP7	T.....T.....	150
WP2	T.....T.....	150
WP5	T.....T.....	150
WP9	T.....T.....	150
Chen	CTTGAATAGTACGACATGGAATAAAGATAGGACTTTAGTTCTATTTTGTGGTTCTAGGACTGAAGTAATGATTAAATAGGACAGTTGGGGCATTGCTATTTAACTGTCAGAGGTGAAATCTTAGATTCTTAAGACGAACTAC	300
WP1		300
WP8		300
WP10		300
WP4		300
WP3	T.....T.....	300
WP6	T.....T.....	300
WP7	T.....T.....	300
WP2	A.....A.....	300
WP5	A.....A.....	300
WP9	A.....A.....	300
Chen	TGCGAAGGCAITTCGAAGGATGTTTTCATTAAATCAAGACGACAGTAGGGGGTTTGAAGACGATTAGATACCGTCGTAATCTCTACCATAACTATGCCGACTAGAGATAGGGAAGCGCT	422
WP1		422
WP8		422
WP10		422
WP4		422
WP3	T.....T.....	422
WP6	T.....T.....	422
WP7	T.....T.....	422
WP2	C.....C.....	422
WP5	C.....C.....	422
WP9	C.....C.....	422

Figure 2 Sequence comparison of 18S rDNA from *Wenyonella philiplevinei* in ducks.




Figure 3 Phylogenetic relationship among the examined cestode species inferred by neighbour joining (NJ), maximum likelihood (ML) and maximum parsimony (MP) analyses based on 18S rDNA sequences, using *Cryptosporidium parvum* as outgroup.

Phylogenetic relationships among the examined *W. philiplevinei* were inferred based on 18S rDNA sequences. Topologies of all trees constructed by different methods (NJ, MP, and ML) with different distance models were similar, with only slight difference of bootstrap values (Fig 3). These results indicate that all the isolates in Hunan province represent *W. philiplevinei*. From the phylogenetic tree, genus *Eimeria* were sister to the *Cyclospora*, consistent with that of previous study (Lopez et al., 1999). The *Wenyonella* were more closely related to the *Eimeria* + *Cyclospora* than to the *Isospora*, with strong support in all of the three phylogenetic analyses (Fig 3).

In conclusion, the 18S rDNA sequences of the *W. philiplevinei* were characterized in ducks from China. The results of the present study have implications for the identification of *W. philiplevinei* infections in ducks in China and elsewhere.

Acknowledgements

This work was supported by the Hunan Province Science & Technology Project (grant no. 2011NK3085) and the Scientific Research Fund of Hunan Provincial Education Department (grant no. 11C0290).

References

- Carlson-Bremer D, Johnson CK, Miller RH, Gulland FM, Conrad PA, Wasmuth JD, Colegrove KM and Grigg ME 2012. Identification of two novel coccidian species shed by California sea lions (*Zalophus californianus*). *J Parasitol.* 98: 347-354.
- Chen JL, Li GQ, Xu QM, Yue CY, Gao ZY, Zhu HB and Liu X 2010. PCR amplification, cloning and sequencing of partial 18S rDNA from *Wenyonella philiplevinei*. *Chin Anim Husb Vet Med.* 37: 67-70.
- Edgar SA 1987. Coccidiosis of ducks and geese. *Vet Hum Toxicol.* 29: 68-70.
- Felsenstein J 1995. PHYLIP (Phylogeny Inference Package), version 3.57c, Department of Genetics, University of Washington, Seattle, WA.
- Huang P, Dong H, Shen J, Han HY and Zhao QP 2004. A preliminary investigation of the coccidian species of ducks in China. *Chin J Prev Vet Med.* 26: 74-77.
- Lopez FA, Manglicmot J, Schmidt TM, Yeh C, Smith HV and Relman DA 1999. Molecular characterization of *Cyclospora*-like organisms from baboons. *J Infect Dis.* 179: 670-676.
- Ogedengbe JD, Hanner RH and Barta JR 2011. DNA barcoding identifies *Eimeria* species and contributes to the phylogenetics of coccidian parasites (Eimeriorina, Apicomplexa, Alveolata). *Int J Parasitol.* 41: 843-850.
- Page RD 1996. TREEVIEW: An application to display phylogenetic trees on personal computers. *Comput Appl Biosci.* 12: 357-358.
- Skírnisson K 1997. Mortality associated with renal and intestinal coccidiosis in juvenile eiders in Iceland. *Parasitologia.* 39: 325-330.
- Strimmer K and Haeseler AV 1996. Quartet puzzling: A quartet maximum likelihood method for reconstructing tree topologies. *Mol Biol Evolution.* 13: 964-969.
- Tamura K, Dudley J, Nei M and Kumar S 2007. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol.* 24: 1596-1599.
- Tewari AK and Maharana BR 2011. Control of poultry coccidiosis: Changing trends. *J Parasit Dis.* 35: 10-17.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F and Higgins DG 1997. The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24: 4876-4882.
- Wünschmann A, Wellehan JF Jr, Armien A, Bemrick WJ, Barnes D, Averbek GA, Roback R, Schwabenlander M, D'Almeida E, Joki R, Childress AL, Cortinas R, Gardiner CH and Greiner EC 2010. Renal infection by a new coccidian genus in big brown bats (*Eptesicus fuscus*). *J Parasitol.* 96: 178-183.
- Yin PY, Jiang JS and Kong FY 1983. Coccidiosis in ducks. *Chin J Vet Med.* 8: 56-58.
- Zhang LX, Ling CS and Li JZ 1999. Prevalent of coccidian in ducks in Henan province, China. *J Anim Sci Vet Med.* 18: 8-10.

